


HADRON STRUCTURE: THE UNSOLVED PUZZLE

Sebastian Kuhn
Old Dominion University
Norfolk, Virginia


Fundamental Problem of Nuclear and Hadronic Physics

- Nearly all well-known (“visible”) mass in the universe is due to hadronic matter
- Fundamental theory of hadronic matter exists since the 1960’s:
 - Quantum Chromo Dynamics
 - “Colored” quarks (u,d,c,s,t,b) and gluons; Lagrangian
- BUT: knowing the ingredients doesn’t mean we know how to build hadrons and nuclei from them!
 - akin to the question:
 - “Given bricks and mortar, how do you build a house?”
- Four related puzzles:
 - What is the “quark-gluon wave function” of known hadrons?
 - How are hadrons (nucleons) bound into nuclei?
 - Does their quark-gluon wave function change inside a nucleus?
 - How do fast quarks and gluons propagate inside hadronic matter?
 - How do fast quarks and gluons turn back into observable hadrons?





Hadron Structure

- Simple-most (constituent quark) model of nucleons (protons and neutrons)
- ... becomes much more complicated once we consider the full relativistic quantum field theory called QCD
- Effective theories: Quark model, χ PT, sum rules, ...
- and Lattice QCD!



Nuclear Structure

- Even more complicated!
- Effective degrees of freedom: nucleons, mesons, nucleon resonances... augmented by phenomenological NN potentials
- Effective theories: low-energy EFT, χ PT, relativistic and non-relativistic potential models, shell model,...
- and Lattice QCD???




How Do We Study Hadron/Nuclear Structure?


- Energy levels: Nuclear and particle (baryon, meson) masses, excitation spectra, excited state decays ->
Spectroscopy (*What exists?*)
- Elastic and inelastic scattering, particle production
Reactions (*Relationships?*)
- Probing the internal structure directly
Imaging (*Shape and Content?*)
- Particular way to encode this: Structure Functions
 - “*Parton wave function*”?
5(6)-dim. Wigner distribution → ...

Overview

- Partonic Structure of the Nucleon
- Polarized and Unpolarized Structure Functions
- Recent Results
 - Spin-Averaged Structure Functions
 - Spin-Dependent Structure Functions
 - Nuclear Structure Functions
- Outlook
 - From 1D to 3D
 - Future Experiments



Parton Distribution Functions



- The 1D world of nucleon/nuclear collinear structure:

- Take a nucleon/nucleus
- Move it real fast along z
 \Rightarrow light cone momentum
 $P_+ = P_0 + P_z (>>M)$
- Select a “parton” (quark, gluon) inside
- Measure **its** l.c. momentum
 $p_+ = p_0 + p_z (m \approx 0)$
- \Rightarrow Momentum Fraction $x = p_+/P_+^*$



- In DIS **: $p_+/P_+ \approx \xi = (q_z - \nu)/M$
 $\approx x_{Bj} = Q^2/2M\nu$

- Probability: $f_1^i(x), i = u, d, s, \dots, G$

In the following, will often write “ $q_i(x)$ ” for $f_1^i(x)$

*) Advantage: Boost-independent along z

***) DIS = “Deep Inelastic (Lepton) Scattering”



Inclusive lepton scattering

Parton model: DIS can access $F_1(x) = \frac{1}{2} \sum_i e_i^2 q_i(x)$ (and $F_2(x) \approx 2xF_1(x)$) Callan-Gross Wandzura-Wilczek

$$g_1(x) = \frac{1}{2} \sum_i e_i^2 \Delta q_i(x) \left(\text{and } g_2(x) \approx -g_1(x) + \int_x^1 \frac{g_1(y)}{y} dy \right)$$

At finite Q^2 : pQCD evolution ($q(x, Q^2), \Delta q(x, Q^2) \Rightarrow$ DGLAP equations), and gluon radiation


$$g_1(x, Q^2)_{pQCD} = \frac{1}{2} \sum_q^{N_f} e_q^2 [(\Delta q + \Delta \bar{q}) \otimes (1 + \frac{\alpha_s(Q^2)}{2\pi} \delta C_q) + \frac{\alpha_s(Q^2)}{2\pi} \Delta G \otimes \frac{\delta C_G}{N_f}]$$

\Rightarrow access to gluons. $\delta C_q, \delta C_G$ - Wilson coefficient functions

SIDIS: Tag the flavor of the struck quark with the leading FS hadron \Rightarrow separate $q_i(x, Q^2), \Delta q_i(x, Q^2)$

Jefferson Lab kinematics: $Q^2 \approx M^2 \Rightarrow$ target mass effects, higher twist contributions and resonance excitations

- Non-zero $R = \frac{F_2}{2xF_1} \left(\frac{4M^2x^2}{Q^2} + 1 \right) - 1, g_2^{HT}(x) = g_2(x) - g_2^{WW}(x)$
- Further Q^2 -dependence (power series in $\frac{1}{Q^n}$)



$$q(x; Q^2), \langle h \cdot H \rangle q(x; Q^2)$$

Traditional "1-D" Parton Distributions (PDFs) (integrated over many variables)


⇒ Our 1D View of the Nucleon

(depends on x and the resolution of the virtual photon $\sim 1/Q^2$)


$$W = \text{final state invariant mass} = \sqrt{M^2 + \left(\frac{1}{x} - 1\right)Q^2}$$

DIS
JLab

- Elastic scattering
(Whole system recoils, $x = 1$, $W = M$)
- Resonances
($x < 1$, $W < 2$ GeV)
- Valence quarks
($x \geq 0.3$, $W > 2$ GeV)
- Sea quarks, gluons
($x < 0.3$)
- “Wee Partons”
($x \rightarrow 0$, Diffraction, Pomerons)



elastic scattering
resonance region
DIS regime: $W > 2$ GeV



Low Q^2 :




ChPT

CQM

OPE

Valence PDFs

- Behavior of PDFs still unknown for $x \rightarrow 1$
 - SU(6): $d/u = 1/2$, $\Delta u/u = 2/3$, $\Delta d/d = -1/3$ for all x
 - Relativistic Quark model: Δu , Δd reduced
 - Hyperfine effect (1-gluon-exchange): Spectator spin 1 suppressed, $d/u \rightarrow 0$, $\Delta u/u \rightarrow 1$, $\Delta d/d \rightarrow -1/3$
 - Helicity conservation: $d/u \rightarrow 1/5$, $\Delta u/u \rightarrow 1$, $\Delta d/d \rightarrow 1$
 - Orbital angular momentum: can explain slower convergence to $\Delta d/d \rightarrow 1$
- Plenty of data on proton \rightarrow mostly constraints on u and Δu
- Knowledge on d limited by lack of free neutron target (nuclear binding effects in d , ^3He)
- Large x requires very high luminosity and resolution; binding effects become dominant uncertainty for the neutron



Moments of Structure Functions

Related to matrix elements of local operators (OPE) - in principle accessible to lattice QCD calculations

Sum rules relate moments to the total spin carried by quarks in the nucleon (and β -decay matrix elements), sea quark asymmetries etc.

At low Q^2 : Higher Twist, Parton-Hadron Duality, Chiral Perturbation Theory, GDH Sum Rule


Bjorken Sum Rule: $\Gamma_1^p - \Gamma_1^n = \frac{g_A}{6} + \text{QCD corr.}$

GDH Sum Rule: $\Gamma_1(Q^2 \rightarrow 0) \rightarrow -\frac{Q^2}{2M^2} \frac{\kappa^2}{4}$

...and γ_0, δ_{LT}

Gottfried Sum Rule:

$$\int_0^1 [F_2^p(x, Q^2) - F_2^n(x, Q^2)] \frac{dx}{x} = \frac{1}{3} - \frac{2}{3} \int_0^1 [\bar{d}(x, Q^2) - \bar{u}(x, Q^2)] dx$$



Unpolarized Structure Functions

