




Higgs Field

- Create “Drag” on Particles (“Molasses”)
- *) Origin of Mass
Makes some gauge bosons very heavy (W 's, Z 's) and therefore short-range (“Weak” interaction)
- Origin of electroweak symmetry breaking
- Pointlike, Fundamental
- Bosons (Spin 0)
- Three massless (“swallowed up” by W 's, Z 's); one very massive (125 GeV)
- Discovered in 2012 at CERN





The LHC at CERN



- See also the movie “Particle Fever”


Higgs Discovery

ATLAS
EXPERIMENT
<http://atlas.ch>





Fundamental Problem of Nuclear and Hadronic Physics

- Nearly all well-known (“visible”) mass in the universe is due to hadronic (=strongly interacting) matter (protons, neutrons, pions, kaons, ...)
- Fundamental theory of hadronic matter exists since the 1960-70’s: **Q**uantum **C**hromo **D**ynamics
 - “Colored” quarks (u,d,c,s,t,b) and gluons; interaction
- BUT: knowing the ingredients doesn’t mean we know how to build hadrons and nuclei from them!
 - akin to the question:
“Given bricks and mortar, how do you build a house?”
- 2 related puzzles:
 - What is the “quark-gluon wave function” of known hadrons?
 - How are hadrons (nucleons) bound into nuclei?
Does their quark-gluon wave function change inside a nucleus?



Hadron Structure

- Simple-most (constituent quark) model of nucleons (protons and neutrons)
- ... becomes much more complicated once we consider the full relativistic quantum field theory called QCD



QCD = Quantum Chromo Dynamics = theory of strong interactions between quarks and gluons

⇒ Our 1D View of the Nucleon


(depends on energy ν and wave length of the virtual photon $\sim 1/Q^2$)

$$W = \text{final state invariant mass} = \sqrt{M^2 + 2M\nu - Q^2}$$


$$x = \text{energy fraction of hit object} = Q^2/2M\nu$$

part


• Elastic scattering
(Whole system recoils, $x = 1$, $W = M$)




• Resonances
($x < 1$, $W < 2 \text{ GeV}$)




• Valence quarks
($x \geq 0.3$, $W > 2 \text{ GeV}$)



• Sea quarks, gluons
($x < 0.3$)




• “Wee Partons”
($x \rightarrow 0$, Diffraction, Pomerons)




S
I
D

elastic scattering resonance region DIS regime: $W > 2 \text{ GeV}$



The Future: 3D Partonic Structure (Holography)



3-D Picture of parton flavor, spin and momentum (TMDs)



- From 1-D to 3-D:




$$q(H, \vec{S}_\perp, x, \vec{k}_\perp, h, \vec{s}_\perp; Q^2)$$




$$q(x; Q^2), \langle h \cdot H \rangle q(x; Q^2)$$

Traditional "1-D" Parton Distributions (PDFs)
(inclusive, integrated over many variables)





$$q(\vec{S}, x, \vec{r}_\perp, \vec{s}; Q^2)$$



3-D parton orbits (GPDs)

Nuclear Structure

- Even more complicated!
- Nuclei effectively look like a bunch of nucleons, mesons, nucleon resonances...
bound together by the strong interaction
- Ultimately, must be explained in terms of quarks and gluons, as well!
- Quark structure might be modified (EMC effect) and in turn affects nuclear binding



Other particle physics puzzles


- What's up with neutrinos?
- What is dark matter?
- What is dark energy?
- Where does inflation come from?
- Are there even more fundamental entities than quarks and leptons?
- Are there unknown forces?

Neutrinos DISAPPEAR!

- Originally discovered by Ray Davis: there are too few neutrinos coming from the sun


- Original experiment in Homestead Mine (Cl): Only 1/3 of expected flux
- Confirmed by Sage, Gallex, Super-K, SNO, ...
- Confirmed with reactors: Bugey, Chooz, KamLand,...
- Also found disappearance of μ -neutrinos in atmosphere: Super-K. Confirmed with K2K

Total Rates: Standard Model vs. Experiment
Bahcall-Pinsonneault 2000



Kamiokande, Super-K



- Detect neutrinos from sun and atmospheric neutrinos
- Only 50% of solar ν 's
- Detection via Cherenkov Light



Kamioka Observatory, ICRR (Institute for Cosmic Ray Research), The University of Tokyo

SNO

- Heavy Water Cherenkov detector
- Sensitive to all 3 types of ν 's with different observables:
 - $d + \nu_e \rightarrow p + p + e^-$;
 - $d + \nu_\mu \rightarrow p + n + \nu_\mu$
- First unambiguous confirmation that total number of ν 's from sun is as expected - only flavor changes





The Nobel Prize in Physics 2015
Takaaki Kajita, Arthur B. McDonald

Share this: [f](#) [G+](#) [t](#) [+](#) 1.5K [✉](#)

The Nobel Prize in Physics 2015



Photo © Takaaki Kajita

Takaaki Kajita

Prize share: 1/2



Photo: K. MacFarlane.
Queen's University
/SNOLAB

Arthur B. McDonald

Prize share: 1/2

The Nobel Prize in Physics 2015 was awarded jointly to Takaaki Kajita and Arthur B. McDonald *"for the discovery of neutrino oscillations, which shows that neutrinos have mass"*


Deficiencies of the Standard Model

The Standard Model is really successful, but...

- Does (fundamental particle rest) mass really come from the Higgs field?
 - Why are the masses so vastly different?
Lowest mass neutrino eigenstate $\nu_1 \rightarrow < 0.01$ eV, highest mass quark t (top quark) $\rightarrow 170,000,000,000$ eV
- Why are there so many “fundamental” particles? (6 leptons, 6 quarks, 1+3+8+1 gauge bosons, Higgs,...). Or are there even more???
- Why are interactions so different in strength? (Gravitation is feeble compared to electroweak and strong interactions)
- ALL IN ALL, why are there so many parameters? (12 fermion masses, 8 mixing angles, 4+1+1 interaction parameters,...) And why are they so finely tuned to allow ordinary matter to exist in our Universe?
- What is the dark matter and dark energy observed in the Universe?
- How can we reconcile gravity with quantum field theory and the other 3 interactions?

Deficiencies of the Standard Model

We observe much more gravitation in the Universe than can be explained by visible mass (and even by all hadronic and leptonic mass left over from the big bang) → WIMPs.




Deficiencies of the Standard Model

Gravitation - what happens at the Planck Scale?

- The Planck Scale - a universal size, time and energy scale
 - Einstein: $E^2 = m^2c^4 + p^2c^2 \Rightarrow E \geq pc$
 - Heisenberg: $\Delta p \cdot \Delta x \geq \hbar/2 \Rightarrow E \geq pc \geq \hbar c/2\Delta x$
 - Newton: $U_{\text{grav}} = m GM/r \Rightarrow$ Escape velocity $v_{\text{esc}} = (2GM/r)^{1/2} \leq c \Rightarrow$
Black hole: Schwarzschild radius $R = 2GM/c^2$
 - Einstein: $M \leftarrow E/c^2 \Rightarrow R = 2GE/c^4 \geq 2G\hbar/(2c^3 R)$
 - \Rightarrow Planck length: $R = (G\hbar/c^3)^{1/2} = 1.6 \cdot 10^{-35}$ m;
Planck Mass $22\mu\text{g}$ (10^{19} GeV)
Planck Energy $2 \cdot 10^9$ J

- What happens at the Planck Scale?

- Space-Time becomes “frothy”
- Pointlike interactions make no sense
- Pointlike particles make no sense



Supersymmetry


- Fundamental Space-Time-Spin symmetry
- Every Particle has a Super-Partner of different spin (different statistics!):
 - Fermions ($S = 1/2$) \Leftrightarrow sFermions ($S = 0$)
 - sneutrinos, selectrons, smus, staus, squarks
 - Bosons ($S = 0,1,2$) \Leftrightarrow Bosinos ($S = 1/2$)
 - winos, zino, photino, gluino, gravitino, higgsino
- May explain dark matter (WIMPs = lightest Super-partner)
- Supersymmetry is broken at high energy scale (1 TeV?) - should be accessible at LHC

Supersymmetry - some (minor?) problems

- Now we are supposed to **double** the number of particles (not a single one has been detected yet)? First LHC run came up empty!
- Add to that a whole bunch of other parameters and possibly new interactions (sfermion decays, quark decays \rightarrow proton should be unstable, but so far only upper limits have been found)
- Why is supersymmetry broken, and why is it broken at yet another mass scale?





Super-Strings

- All particles are vibrations of incredibly tiny strings (of size of the Planck scale, 10^{17} times smaller than resolution of present accelerators). Tension = $10^9\text{J}/10^{-35}\text{m} = 10^{40}$ tons
- They are “wrapped” around extra dimensions
- Their vibrational energies determine their masses.
- Vibration patterns determine charges and spin (determined by geometry of extra dimensions).
- Original idea: Kaluza-Klein.




Super-Strings

- Require 9+1 dimensions to avoid negative probabilities
- Extra dimensions “curled up”
- “Calabi Yau Spaces”
- Compare to ants on a hose



Super-String Theory

- Unified picture of all four interactions
- Avoids singularities in particle interactions - you can't make them smaller than the Planck Length
- Includes Supersymmetry "automatically"
- Could be compatible with all 4 forces uniting in strength at the Planck scale
- Might explain beginning of Universe



Super-Strings - some (minor?) Problems

- Nobody can write down the exact theory (equations aren't fully known)
- Only approximate solutions known
- Many competing versions (Brane theory...) -> too many solutions
- Presently hard to see how we can test them experimentally

Nobel Laureate 2004



The New York Times
nytimes.com

PRINTER-FRIENDLY FORMAT
SPONSORED BY STARBUCKS.COM

September 2, 2003

One Cosmic Question, Too Many Answers

By DENNIS OVERBYE

Call it the theory of anything.

But the same calculations confirmed that string theory could have a vast number of solutions, each representing a different universe with slightly different laws of physics. The detailed characteristics of any particular one of these universes — the laws that describe the basic forces and particles — might be decided by chance.

As a result, string theorists and cosmologists are confronted with what Dr. Leonard Susskind of Stanford has called "the cosmic landscape," a sort of metarealm of space-times. Contrary to Einstein's hopes, it may be that neither God nor physics chooses among these possibilities, Dr. Susskind contends. Rather it could be life.

Only a fraction of the universes in this metarealm would have the lucky blend of properties suitable for life, Dr. Susskind explained. It should be no surprise that we find ourselves in one of these. "We live where we can live," he said.


Dr. Susskind conceded that many colleagues who harbor the Einsteinian dream of predicting everything are appalled by that notion that God plays dice with the laws of physics.

Among them is **Dr. David Gross**, director of the Kavli Institute of Theoretical Physics in Santa Barbara, Calif., who said, "I'm a total Einsteinian with respect to the ultimate goal of science." Physicists should be able to predict all the parameters of nature, Dr. Gross said, adding, "They're not adjustable."

The Future ?



- LHC (CERN)



- NLC ?