

HAUSTORIUM

Parasitic Plants Newsletter

Official Organ of the International Parasitic
Seed Plant Research Group



May 1994

Number 29

● ● SIXTH INTERNATIONAL PARASITIC WEED SYMPOSIUM 1996

The next symposium is tentatively scheduled for **1996** in Cordoba, Spain. If you are interested in attending, fill out the enclosed insert and mail directly to the address on the insert. The first circular is scheduled for mailing in September **1993**. For further information contact:

Dr Maria Teresa Moreno
Centro de Investigacion y Desarrollo Agrario
Apartado **4240**
14080 Cordoba
SPAIN
Fax: (country code **34**) **57 202721**, Telex: **76686**

● THIRD INTERNATIONAL WORKSHOP ON OROBANCHE

Held in Amsterdam **8-12** November **1993**, the workshop had about **100** delegates from **23** countries who enjoyed an excellent third Orobanche workshop in the impressive surroundings of the Royal Tropical Institute. Sixty papers and 30 posters covered many aspects of both Orobanche and Striga. After introductory papers contributions on the basic biology and taxonomy of Orobanche included observations on the evolution of the Orobanchaceae as deduced from the plastid genome (Thalouarn). There was repeated discussion of the taxonomic status of *O. cernua* v. *O. cumana* and *O. ramosa* v. *O. aegyptiaca* but no resolution. Techniques based on pollen morphology (Abu Sbaiah) and chemotaxonomy (Andary) had been used to confirm distinctions between the sections *Trinychon* and *Orobanche* (*Osproleon*) of the genus but not for separating the species within them. The conditioning and germination processes of *O. aegyptiaca* were described in new detail (Mayer); a new in vitro tech-

nique for growing host roots should throw useful light on stimulant exudation (Croes); and the importance of primary and secondary dormancy was discussed (Piet-erse). There were also new studies on the precise role of ethylene in *Striga* germination (Babiker; Fer; Thuring). For the latter study an elegant new ethylene-detection technique had been used. Ethylene is not apparently involved in *Orobanche* germination (Joel; Wegmann). *Strigol* has now been confirmed as the natural *Striga*-germinating stimulant from maize and proso millet (Butler). The *Orobanche* stimulants have still not been fully determined but are increasingly thought also to be members of the *strigol* family (Wegmann). Two groups have continued work on structure-activity relations of *strigol* analogues (Zwan-wenburg; Welzel). Haustorial initiation in *Orobanche* remains to be explained but the penetration process into the host was well described (Joel; Doerr; Shomer-Elan; Losner-Goshen). Progress in host crop resistance was reported for sunflower. Workers in Spain (Melero-Vara), Bulgaria (Encheva) and Turkey (Pet-zoldt) reported development of immunity to all major pathotypes of '*O. cumana*' but further pathotypes were expected to develop and there was also a report from Israel of failure of resistance under low temperature conditions (Jacobsohn). There was reiteration of the hypothesis that resistance results from a lignification response in the pericycle (Antonova) but also an alternative suggestion that resistance is expressed primarily by development of an isolation layer in the cortex and that lignification is a later secondary effect (Doerr). The possible role of phytoalexins in resistance was discussed (Wegmann). Breeding and selection in faba bean has yielded lines with useful levels of resistance to *O. crenata*, often combined with relative tolerance, for use in Egypt (Zaitoun; Abdalla; Khalil), and in Spain (Cubero); and with resistance to *O. foetida* in Tunisia (Kharrat). Among other control measures, new imidazolinone herbicides, especially imazethapyr and imazaquin, show promise not only in

faba bean but also in other legumes such as pea and perhaps in sunflower (Garcia-Torres; Saber; Jacobsohn; Castejon-Munoz). In discussion, however, there was a warning that resistance to this class of herbicides is likely to develop very rapidly (Gressel). Maleic hydrazide is still being used successfully in tobacco in Cuba (Labrada). Mechanical control by 'mini-spear' was again endorsed for use in tobacco (Krishna Murty). There were further encouraging indications of the potential for mycoherbicides based on *Fusarium* species from China and from Bulgaria and India (Bozoukov; Bedi); also a report of successful biocontrol of *Striga hermonthica* by *F. nygamai* (Sauerborn). Integrated systems involving rotation, delayed planting etc had been tested in Morocco, Syria and Spain (Kroschel; Schnell; Solh; Lopez-Granados) and useful information obtained on the fate of the seed bank under different regimes. Where fertilizer was being included it was generally assumed to be increasing crop tolerance rather than contributing directly to control. Reports of Orobanche problems and current research in different countries included papers from Chile, Cuba and Sudan, where *O. ramosa* is a problem in all three countries (Kogan; Labrada; Babiker), also from Bulgaria, Spain, India, Morocco, Tunisia, Egypt, Jordan and Israel. Reports on EC projects included several on *Striga*, from France (Salle; Raynal-Roques; Pari), Benin (Gbehounou), U.K. (Lane), Kenya (Ransom; Odhiambo; Baltus; Kuiper) and Netherlands (Verkleij). Ransom presented the first encouraging results from a longterm trial in which the effects of hand-pulling and other treatments were being monitored for their effects on the *S. hermonthica* seed bank. In a final session of general papers, Dr Gressel presented an authoritative account of the potential for the application of genetic engineering to the problem of parasitic weeds, especially the development of herbicide-resistant crops. The potential had already been proven with crops engineered for resistance to sulphonyl urea and imidazolinone herbicides but he warned that resistance to these two groups is likely to break down rapidly, while that to glyphosate should be more durable. Chris Parker summarized the latest prospects for control and emphasized the need for integrated systems which eliminated or at least minimized the risks of breakdown of crop resistance and/or the development of herbicide resistance in the parasite. Dr Abu-Irmaileh reported on the conclusions from the FAO Orobanche meeting held in Amman, Jordan in October and emphasized the need to reduce the risks of new infestations which arose from careless use of unfermented manure and contaminated crop seed. Dr Labrada also outlined a new FAO proposal for a coor-

dated Orobanche research programme in the Near East Region. Finally Dr Pieterse announced plans for the creation of a research group on Parasitic Weeds to be set up in the framework of the European Weed Research Society (EWRS) (regrettably since rejected by EWRS - Ed.). Dr Pieterse and his colleagues are to be congratulated on the excellent facilities and organization provided for this workshop. The proceedings are expected to be available in a few months time. Details will be in the next issue of HAUSTORIUM or contact A. H. Pieterse, Royal Tropical Institute, Mauritskade 63, 1092 ED, Amsterdam, The Netherlands.

● **PYRULARIA PUBERA AS A PATHOGEN OF CHRISTMAS TREES IN WEST VIRGINIA; THESIUM IN MONTANA**

The Santalaceae is a small family in North America with five genera each with a single species. Although they are all root parasites, there has been no recorded host damage. A sixth genus, *Thesium*, was reported as a minor weed in the Great Plains several decades ago but has since been extirpated. Recently a species *Thesium* and a native shrub of the Santalaceae have been reported as weeds. During the 1993 growing season, the West Virginia forest authorities were notified of damage to *Abies*, *Picea*, and *Pinus* which are grown in plantations in the Appalachian Mountains of southern West Virginia as Christmas trees. Trees were losing leaves, stems were dying back, and in heavy infestations trees were killed. Examination of the roots revealed white swellings which were assumed to be nematodes but were later determined to be haustoria of an unknown parasitic plant. During a site visit in May, it was determined that the causative agent is the widespread and common Appalachian shrub, *Pyrrularia pubera*, known locally as buffalo nut. This is the first report of this genus causing damage anywhere. Control measures are being developed but the strongly rhizomatous nature of the parasite makes control difficult. The genus *Thesium* has been introduced into the United States on several occasions and a new infestation has been discovered on range land in Montana. The tentative determination of this species is *Thesium humile*, a documented problem in some grain crops in the Middle East, but the Montana plant remains to be verified as the genus is a large one with considerable taxonomic difficulty.

S. Clark Haynes, West Virginia Department of Agriculture, Charleston, West Virginia 25305-0170 and L. J. Musselman.

● **AN IMPORTANT NEW BOOK ON PARASITIC PLANTS**

Parasitic Weeds of the World: Biology and Control. C. Parker and C. R. Riches, 1993. 332 pages, 16 plates + 44 Figures. ISBN 0 85198 873 3. Price 45 pounds sterling. Available from CAB International Wallingford, Oxon OX10 8DE, UK; North America: 845 North Park Avenue, Tucson, AZ 85719, USA. Price US \$89.50. Copies may be available to developing country institutions at a reduced cost from Technical Centre for Agricultural and Rural Cooperation, P.O. Box 380, 6700 AJ Wageningen, Netherlands. The book includes 16 colorful high quality plates. In sharp contrast to the plates, most of the nearly 44 line drawings fall short of textbook standard. Still, I found them very helpful in illustrating the general features and morphology. It is divided into six chapters: 1- *Striga* (9 spp); 2- *Striga gesnerioides* and *Alectra* (3 spp); 3- *Orobanchae* (6 spp); 4- Other root parasites in Scrophulariaceae, Orobanchaceae, Santalaceae, and Balanophoraceae; 5- *Cuscuta* (14 spp) and *Cassytha*; 6- the Loranthaceae and the Viscaceae. There are keys to species and each species is discussed separately as to its distribution, host range, variation, economic importance, biology and physiology, ecology, and control. The aspect of control has an interesting part to it- recommending different control measures for various levels of parasitic infestation. In addition, each chapter concludes with specific control strategy appropriate to the parasite stressing integrated control measures and improved extension services. The book draws from and integrates the vast literature on the various aspects of parasitic plants that have accumulated in the past 25 years (850 references). Although not in great depth, it is one of the few publications that bring together all the major groups of parasitic plants and address the different aspects of their biology, ecology, physiology, and control. A comprehensive and up-to-date review of the control measures of the economically important parasitic weeds make this book special. Thus it will serve as one of the major sources of information to students and research workers.

Kamal I. Mohamed, Old Dominion University, Norfolk, Virginia.

● **STRIGA IN COTE D'IVOIRE 1992 AND**

A reconnaissance was undertaken in early September 1992 of rice areas north of Boundiali, Korhogo and Ferkessedougou. In the west of this zone *S. aspera* is conspicuous among wild grasses and locally as severe infestations in rice and maize. It is also common on "fonio" (*Digitaria exilis*). *Striga hermonthica* occurs at low densities in rice and maize and associated also with sorghum and grass weeds, especially *Digitaria horizontalis*. In the vicinity of Korhogo and Ferkessedougou where *S. aspera* appears not to occur, severe infestations of *S. hermonthica* were seen in rice, maize, and sorghum. Patches of the yellow flowered morphotype of *S. asiatica* were also seen on rice. Further south in the forest/savannah transition zone, we found both red and yellow morphotypes of *S. asiatica* parasitic on wild grasses. Farmers had not seen this parasite on rice. *Striga gesnerioides* was observed on *Ipomoea eriocarpa* near Korhogo. No *Striga* was apparent in cowpea and farmers were not aware of the parasite on this crop so it seems likely that this area is south of the range of the cowpea strain of *S. gesnerioides*. *Alectra vogellii*, on the other hand, is common on peanut in the Korhogo areas with noticeable damage at several sites, The Natural Resources Institute (NRI) is collaborating with the West African Rice Development Association (WARDA) in a search for *Striga* resistance for use in rice varieties suitable for savannah areas of West Africa.

C. R. Riches, D. E. Johnson, NRI, UK and R. Diallo, WARDA, Bouake

The results of our survey in mid-October 1993 are quite different from those reported above likely due to a later season. We found only one field of sorghum and maize severely infested with *S. hermonthica*, about 70 km west of Ferkessedougou. *Striga brachycalyx* is abundant in native grasslands. Several other species were found but no *S. aspera*.

D. M. Berner, International Institute of Tropical Agriculture and L. J. Musselman

● **LITERATURE**

Aparicio, A. 1993. Sex-determining and floating translocation complexes in *Viscum cruciatum* Sieber ex Boiss. (Viscaceae) in southern Spain. Some

- evolutionary and ecological comments. Botanical Journal of the Linnean Society **111**: 359-369.
- Appa Rao, S., A.W. Mukred, M.H. Menagesha, H.M. Amer., K.N. Reddy, A. Alsurai and M. Saeed. **1993**. Collecting crop germplasm in Yemen. FAO/IBPGR Plant Genetic Resources Newsletter **94/95**: 28-31. (Notes the importance of both *Striga hermonthica* and *S. asiatica* on sorghum and pearl millet in Yemen).
- Babiker, A. G. T., N. E. Ahmed, A. H. Mohamed, M. E. El Mana and S. M. E. Tayeb. **1993**. *Striga hermonthica* on sorghum: chemical and cultural control. In: Proceedings Brighton Crop Protection Conference Weeds - **1993**, pp. 907-911.
- Baenziger, H., V. Maschwitz and G. Zizka. **1993**. Rafflesiaceae-eine faszinierende und raetselhafte pflanzenfamilie. Der Palmengarten (Munich) **2/93**: 65-74. (A beautifully illustrated account of the Rafflesiaceae including full color pictures of *Rafflesia* and holoparasites of some other families).
- Baumel, P., W. D. Jeschke, L. Witte, F. C. Czygan and P. Proksch. **1993**. Zeitschrift fur Naturforschung, Section C, Biosciences **48**: 436-443.
- Barlow, B. A. and R. Schodde. Bird dispersal of Loranthaceous mistletoes to remote Pacific islands: Symbiosis in default. Eeaufortia **43(8)**: 124-129. (Suggests that once distance dispersal by birds of mistletoes to remote islands occurs, the species may be freed from constraints imposed by avian dispersal so range extension may be increased. There may be a self generated distance dispersal syndrome for the Pacific which has arisen by default).
- Bergmann, C., K. Wegmann, K. Frischmuth, E. Sampson, A. Krantz, D. Welgelt, P. Koll and P. Welzel. **1993**. Stimulation of *Orobanche crenata* seed germination by (+)-strigol and structural analogues: dependence on constitution and configuration of the germination stimulants. Journal of Plant Physiology **142**: 338-342.
- Berner, D. K., A. E. Awad and E. I. Aigbokhan. **1992**. Efficacy of imazaquin seed treatment for the control of *Striga gesnerioides* and *Alectra vogelii* in cowpea (*Vigna unguiculata*). (abstract) Phytopathology **82**: 1108. (Treating cowpea seed with imazaquin - 5 minute soak in 0.33% solution of ammonium salt - prevented emergence of both parasites and increased cowpea biomass by 73%).
- Berner, D. K., K. F. Cardwell, B. O. Faturoti, F. O. Ikie and O. A. Williams. **1994**. Relative roles of wind, crop seeds and cattle in dispersal of *Striga* spp.. Plant Disease **78**: 402-406. (Concludes crop seed more important than either wind or cattle).
- Boemmer, D., G. Haberhausen and K. Zetsche. **1993**. A large deletion in the plastid DNA of the holoparasitic flowering plant *Cuscuta reflexa* concerning two ribosomal proteins (rpl2/rpl23), one transfer RNA (trnI) and an ORF 2280 homologue. Current Genetics **24**: 171-176.
- Boussim, I. J., G. Salli and S. Guinko. **1993**. *Tupinanthus* parasite du kariti au Burkina Faso. Part 1: Identification et distribution; Part 2: Phnologie, biologie et digbts. Bois et Forets des Tropiques **238 (4)**: 45-52 and 53-65. (Informative and well-illustrated description of the problems caused by *Tapinanthus dodoneifolius*, *T. globiferus* and *T. ophioides*. In French but with an extended English summary).
- Carson, A. G. **1993**. Studies on bio and chemical control of *Striga hermonthica* (Del.) Benth. in the Gambia. Nuisibles-Pests-Pragas **1**: 71-81. (Reporting 80-90% suppression of *Striga* by *Cassia obtusifolia* and a small increase in maize yield when removed after 10 days).
- Castejon-Muqoz, M., L. Garcia-Torres and F. Lopez-Granados. **1993**. [Chronology of *Orobanche crenata* infestation in sunflower in Andalucia (southern Spain) and present problem] (in Spanish). Actas, Congresso **1993**, Sociedad Espanola de Malherbologia, Lugo, **1993**, pp. 154-157.
- Chitere, P. O. and B. A. Omolo. **1993**. Farmers' indigenous knowledge of crop pests and their damage in western Kenya. International Journal of Pest Management **39**: 126-132. (Over 40% of farmers in W. Kenya reported *Striga* occurrence).
- Clark, L. J., K. G. Shawe, G. Hoffman and G. R. Stewart. **1994**. The effect of *Striga hennonthica* (Del.) Benth. infection on gas-exchange characteristics and yield of a sorghum host, measured in the field in Mali. Journal of Experimental Botany **45**: 281-283. (This study failed to confirm any specific reduction of host photosynthesis by *Striga* infection, suggesting that previous reports of reduced photosynthesis could be due to an artifact of pot experimentation).
- Cordero, C. M., A. M. G. Serrano and M. J. A. Gonzalez. **1993**. Transfer of bipiperidyl and quinolizidin alkaloids to *Viscum cruciatum* Sieber (Loranthaceae) hemiparasitic on *Retama sphaerocarpa* Bossier (Leguminosae). Journal of Chemical Ecology **19**: 2389-2393.
- Cubero, J. I., A. H. Pieterse, S. A. Khalil and J. Sauerborn. **1993/1994**. Screening techniques and sources of resistance to parasitic angios-

- perms. *Euphytica* 73 (1-2): 51-58. (A review).
- Cudney, D. W., S. B. Orloff and D. A. Demason. 1993. Effects of thiazopyr and trifluralin on dodder (*Cuscuta znedcora*) in alfalfa (*Medicago sativa*). *Weed Technology* 7: 860-864.
- Danko, J. 1993. Use of herbicide Roundup for control of broomrape (*Orobancha ramosa* L.) on tobacco. *Ochrana Rostlin (Plant Protection) (Prague)* 29: 303-307. (in Slovak) (0.0175% or 0.035% twice at 55-60 days + 14-20 days later, applied to lowest 15 cm, gave up to 90% control but some damage to Virginia tobacco).
- Dizdaroglu, T., G. Eginlioglu and C. O. Sabanci. 1993. (The main technical and economic problems of alfalfa production in Aegian and South Marmara regions). *Anadolu* 3: 11-40, (in Turkish) (13% of the crop area is affected by *Cuscuta* spp).
- DeJongh, K., P. van Damme and D. C. Berner. 1993. Suicidal germination of *Striga* seed using several trap crop varieties. *Mededelingen van de Faculteit van de Landbouwweten schappen, Universiteit Gent* 58/3a: 983-993.
- Doyle, G. J. 1993. *Cuscuta epithymum* (L.) L. (Convolvulaceae), its hosts and associated vegetation in a limestone pavement habitat in the Burren Lowlands in County Clare (H9), Western Ireland. *Biology and Environment, Proceedings of the Royal Irish Academy, Section B* 93B: 61-67. (Although attacking a wide range of hosts, *C. epithymum* may depend on a narrow range of primary hosts for initial establishment).
- Fahmy, G. M. 1993. Transpiration and dry matter allocation in the angiosperm root parasite *Cynomorium coccineum* L. and two of its halophytic hosts. *Biologia Plantarum* 35: 603-608. [Transpiration of *C. coccineum* was lower than that of its hosts (as in Ouf, 1993). Infection reduced transpiration of the hosts.]
- Flueckiger, H. (+ 8 co-authors). 1993. Mistletoe through the seasons. 1994 Calendar. Weleda AG Pharmaceuticals, Arlesheim, Switzerland. (An unusual calendar with illustrations of *Viscum album* and very informative text concerning the biology, mythology and medicinal uses of *V. album*).
- Garcia Torres, L. 1993. *Biología y Control de Especies Parasitas*. Editorial Agrícola Espanola. 94 pages. (This well illustrated booklet contains a great deal of information on the biology and control of all major parasitic plants. It is different from numerous other books and booklets on the topic which have appeared in the past few years in that it contains many helpful graphs. The booklet is in Spanish but deserves translation into English for a wider distribution).
- Gibson, W. 1993. Selective advantages to hemiparasitic annuals, genus *Melampyrum*, of a seed-dispersal mutualism involving ants: II Seed-predator avoidance. *Oikos* 67: 345-350. (Ants shown to have less effect than expected in deterring rodent seed predators).
- Hawksworth, F.G. and Wiens, D. 1993. Change in status of a dwarf mistletoe (*Arceuthobium*, Viscaceae) from China. *Novon* 3: 156. (*Arceuthobium pini* var. *sichuanense*, parasitising *Picea* spp. raised to specific rank as *A. sichuanense*).
- Heide-Jorgensen and J. Kuijt. 1993. Epidermal derivatives as xylem elements and transfer cells: a study of the host-parasite interface in two species of *Triphysaria* (Scrophulariaceae). *Protoplasma* 174: 173-183. (Further contributions to understanding the remarkable versatility of the haustorium. In these species of *Triphysaria*, formerly *Orthocarpus*, the epidermal cells develop into xylem elements and transfer cells).
- Hezewijk, M van, 1994. Germination Ecology of *Orobancha crenata* - Implications for Cultural Control Measures. Doctoral Thesis, Vrije Universiteit, Amsterdam. 162 pp. (Detailed studies of conditioning and germination of the seeds of *O. crenata* and the effects of temperature, sowing date and nitrogen on field infestations).
- Hezewijk, M. J. van, A. P. van Beem and J. A. C. Verkleij. 1993. Germination of *Orobancha crenata* seeds as influenced by conditioning temperature and period. *Canadian Journal of Botany* 71: 786-792. (Optimum temperature for germination was 15-20 ° C and optimum length of conditioning at 20 ° C was 18 days. Conditioning for more than 4 weeks can lead to secondary dormancy).
- Hosmani, M. M., T. D. Malipatil and M. Hanumanthappa. 1993. *Orobancha* and Its Control. ICAR, New Delhi, 31 pp.
- Huang, J. Z., Y. H. Li and Y. Y. Zhang. 1993. (Surface ultra-structure of seeds of Chinese *Cuscuta*). (in Chinese) *Acta Phytotaxonomica Sinica* 31: 261-265.
- Ilan, A. S. 1993. Germinating seeds of the root parasite *Orobancha aegyptiaca* Pers. excrete enzymes with carbohydrase activity. *Symbiosis (Rehovot)* 15:61-70.
- Ish-Shalom-Gordon, N., R. Jacobsohn and Y. Cohen. 1993. Inheritance of resistance to *Orobancha cumana* in sunflower. *Phytopathology* 83:

- 1250-1252. (Resistance of SW501 and RW637 to *O. cumana* in Israel is attributed to a single dominant gene).
- Jia Zhong-Jian and Gao Jian-Jun. 1993. Phenylpropanoid glycosides from *Pedicularis striata* Pall ssp. *arachnoidea*. *Phytochemistry* 34: 1188-1190.
- Jones, B. M. G. 1993. Genetics of resistance to *Striga hermonthica*. In: Rezene Fessehaie (Ed.) Proceedings, 8th Annual Conference of the Ethiopian Weed Science Committee, Addis Abeba, 1990, pp. 1-9.
- Kassahun Zewdie and Tadesse Eshetu. 1993. Preliminary observation on feasibility of mechanical control of parasitic plants on trees. In: Rezene Fessehaie (Ed.), Proceedings, 7th Annual Conference, Ethiopian Weed Science Committee, Addis Abeba, 1989, pp. 41-42.
- Kleifeld, Y., Y. Goldwasser, G. Herzlinger, D. M. Joel, S. Golan and D. Kahana. 1994. The effects of flax (*Linum usitatissimum* L.) and other crops as trap and catch crops for control of Egyptian broomrape (*Orobanche aegyptiaca* Pers.) *Weed Research* 34: 37-44. (Reporting the moderate effectiveness of flax as a rotational cleaning crop).
- Kuijt, J. 1994. A second Brazilian species of *Ixocactus* (Loranthaceae). *Brittonia* 46(1): 72-74.
- Lane, J. A., J. A. Bailey, R. C. Butler and P. J. Terry. 1993. Resistance of cowpea [*Vigna unguiculata* (L.) Walp.] to *Striga gesnerioides* (Willd.) Vatke, a parasitic angiosperm. *New Phytologist* 125: 405-412.
- Lane, J. A., D. V. Child, T. H. M. Moore, H. M. Frost, P. J. Terry and J. A. Bailey. 1993. Host-specificity of *Striga* species and evaluation of cowpea and *Sorghum* germplasm for resistance. In: Proceedings Brighton Crop Protection Conference Weeds - 1993, pp. 937-942.
- Lane, J. A., T. H. M. Moore, D. V. Child, K. F. Cardwell, B. B. Singh and J. A. Bailey. 1994. Virulence characteristics of a new race of the parasitic angiosperm, *Striga gesnerioides*, from Benin on cowpea (*Vigna unguiculata*). *Euphytica* 72: 183-188. (A population of *S. gesnerioides* from S. Benin is the first able to parasitise B.301, but 58-57 and IT81D-994 are resistant to it).
- Lee, R. T., H. J. Gabius and Y. C. Lee. 1992. Ligand binding characteristics of the major mistletoe lectin. *Journal of Biological Chemistry* 267: 23722-23727.
- Linke, K.-H., A. M. Abd El-Moneim and M. C. Saxena. 1993. Variation in resistance of some forage legumes species to *Orobanche crenata* Forsk. *Field Crops Research* 32: 277-285. [Species studied included *Lathyrus ochrus* (resistant), *L. sativus* and *L. cicera* (susceptible), and *Vicia narbonensis*, *V. sativa* and *V. villosa* (showing varietal differences)],
- Lopez-Granados, F. and L. Garcia-Torres. 1993. Population dynamics of crenate broomrape (*Orobanche crenata*) in faba bean (*Vicia faba*). *Weed Science* 41: 563-567. (Infestation of *O. crenata* increased about 3-fold each year with repeated cropping of faba bean).
- Luo, M., G. E. Su, J. H. Liu and T. G. Ao. 1993. (Study on the seed germination of *Cynomorium songaricum* without host). (in Chinese) *Grassland of China*, 1993 (2): 62-64. (Reporting up to 68% germination with various combinations of host root extract, auxin and cytokinin. A low temperature treatment increased germination).
- Ma, C. M., S. S. Jia, T. Sun and Y. W. Zhang. 1993. (Triterpenes and steroidal compounds from *Cynomorium songaricum*). *Acta Pharmaceutica Sinica* 28: 152-155.
- Marwat, K. B., G. R. Marwat and F. M. Sarim. 1993. A checklist and key to parasitic weeds of Pakistan. *Pakistan Journal of Weed Science Research* 6 (1-2): 1-15.
- Mbwaga, A. M., S. Pande, W. A. J. de Milliano and R. I. Karunakar. 1993. Diseases and parasitic weeds of pearl millet in Tanzania: occurrence and incidence from 1986-1990. *Phytophylactica* 25: 271-273. (*Striga hermonthica* and *S. asiatica* of very minor importance in pearl millet, even when adjacent to seriously infested sorghum, maize or finger millet).
- McCormac, J. S. and J. L. Windus. 1993. Fire and *Cuscuta glomerata* Choisy in Ohio: A connection? *Rhodora* 95(882):158-165.
- Mohamed, K. I. 1994. Biosystematics and Diversification of the genus *Striga* Lour. (Scrophulariaceae) in Africa. PhD Dissertation, Old Dominion University. 170 pages.
- Mohamed-Ahmed, A. G. and D. S. H. Drennan. 1993. Bioassay studies on germination of *Orobanche ramosa* with root exudates and extracts. In: Proceedings Brighton Crop Protection Conference Weeds - 1993, pp. 913-918.
- Mumera, L. M. and F. E. Below. 1993. Role of nitrogen in resistance to *Striga* parasitism of maize. *Crop Science* 33: 758-763. (Reporting beneficial effects of three different forms and four timings of nitrogen application in field trials),
- Murphy, S. R., N. Reid, Z. Yan and W. N. Venables. 1993. Differential passage time of mistletoe fruits through the gut of honeyeaters and flower-

- peckers: effects on seedling establishment. *Oecologia* 93: 171-176. [Seeds of *Amyema quandang* germinated over 80% after passage through both birds but establishment after 5 months was better after shorter passage through mistletoe-birds (12 m) v. honeyeaters (40 m)].
- Near East Working Group for Improved Weed Management. 1993. Newsletter 14: 1-8. (Newsletter devoted to account of the Workshop on Parasitic Weeds held in Amman, Jordan in September 1993).
- Okonkwo, C. A. C. and A. Garba. 1993. Selection of *Striga* resistant sorghum germplasm in the northern Guinea Savanna. *Plant Varieties and Seeds* 6: 197-205. (147 varieties tested for resistance/tolerance. Bauchi Local noted as apparently immune).
- Orloff, S. B. and D. W. Cudney. 1993. Controlling dodder in alfalfa hay calls for an integrated procedure. *California Agriculture* 47(6): 32-35. (Suggesting various combinations of trifluralin, mowing, burning and flail mowing).
- Ouf, S. A. 1993. Mycological studies on the angiosperm root parasite *Cynomorium coccineum* L. and two of its halophytic hosts. *Biologia Plantarum* 35: 591-602. (Infection by *C. coccineum* resulted in reduced fungal population on shoots of hosts *Limonium delicatulum* and *Arthrocnemum glaucum*).
- Panvini, A. D. and W. G. Eickmeier. 1993. Nutrient and water relations of the mistletoe *Phoradendron leucarpum* (Viscaceae): how tightly are they integrated? *American Journal of Botany* 80: 872-878.
- Parker, C. 1993. Weed management strategy for parasitic weeds. In: *Integrated Weed Management for Sustainable Agriculture*. Souvenir 1993, from International Symposium, Hisar, 1993. Indian Society of Weed Science. pp. 49-53.
- Parker, C. and C. R. Riches. 1993. *Parasitic Weeds of the World: Biology and Control*. CAB International, Wallingford, U.K. 340 pp. (See review in this issue of HAUSTORIUM).
- Press, M. C. and I. Cechin. 1993. Influences of nitrogen on the interaction between *Striga hermonthica* and its sorghum host: implications for control. In: *Proceedings Brighton Crop Protection Conference Weeds-1993*, pp. 931-936.
- Raynal-Roques, A. 1993. Contribution a la connaissance de la biologie des *Striga* (Scrophulariaceae): types biologiques et phinologie. *Bulletin Musee Nationale de l'Histoire Naturelle, Paris 4e Series* 15, section B. *Adansonia* (1-4): 3-21. [Classifying 39 taxa into rainy season annuals, early dry-season annuals and perennials (18 taxa), describing the means of perennation, and proposing that *S. fulgens* should be excluded from the Scrophulariaceae].
- Reuveni, J., J. Gale and A. M. Mayer. 1993. Reduction of respiration by high ambient CO₂ and the resulting error in measurements of respiration made with O₂ electrodes. *Annals of Botany* 72: 129-131. (Study included *Orobancha aegyptiaca*).
- Saghir, A. R., S. Itani and A. Ahmad. 1993. *Cuscuta* control in vegetable crops in the United Arab Emirates. Near East Working Group for Improved Weed Management Newsletter 13: 2. (Glufosinate-ammonium selectively controlled *C. campestris* in mint at 25 ppm).
- Salatino, A., J. E. Kraus and M. L. F. Salatino. 1993. Contents of tannins and their histological localization in young and adult parts of *Struthanthus vulgaris* Mart. (Loranthaceae). *Annals of Botany* 72: 409-414.
- Salle, G., H. Frochot and C. Andary. 1993. Le gui. (The mistletoe) *La Recherche* 260 (Dec. 1993): 1334-1342. (A valuable, beautifully illustrated review of the biology of *Viscum album*, including many original observations).
- Sanders, I. R., R. T. Koide and D. L. Shumway. 1993. Mycorrhizal stimulation of plant parasites. *Canadian Journal of Botany* 72: 1143-1146. (Successful parasitism of *Abutilon theophrasti* by *Cuscuta pentagona* depended on mycorrhizal infection of host roots).
- Sauerborn, J. 1993. Natural antagonists for parasitic weed control in agroecosystems. *Proceedings, International Symposium, Indian Society of Weed Science, Hisar, 1993, Volume 1*, pp. 29-33.
- Scharpf, R. F. and F. G. Hawksworth. 1993. Mistletoes In: Scharpf, R. F. (ed). *Diseases of Pacific Coast Conifers*. USDA Forest Service Agriculture Handbook 521 (new revised edition), USDA, Washington, 112-135. [A well-illustrated account of the identification and importance of *Phoradendron* (4 spp). and *Arceuthobium* (17 taxa)].
- Seel, W. E. and M. C. Press. 1993. Influence of the host on three sub-Arctic annual facultative root hemiparasites. I. Growth, mineral accumulation and above ground dry-matter partitioning. *New Phytologist* 125: 131-138. (*Rhinanthus minor*, *Euphrasia frigida* and *Melampyrum sylvaticum* all very greatly more vigorous with host than without. *Rhinanthus minor* and *E. frigida* more vigorous on a legume than on a grass).

- Shawe, K. G. and M. J. Ingrouille. **1993**. Isoenzyme analysis demonstrates host selection of parasite pathotypes in the association between cowpea and *Striga gesnerioides*. In: Proceedings Brighton Crop Protection Conference Weeds - **1993**, pp. **919-924**.
- Siddiqui, S. U., S. Khalid, R. A. Shad and K. B. Marwat. **1993**. Host range and distribution of *Orobanche aegyptiaca* and *O. cernua*: two important parasitic weeds of crops in Pakistan. Pakistan Journal of Weed Science Research **6 (1-2): 61-67**. (Important host crops in Pakistan include potato, tomato, tobacco, onion and brassicas).
- Simier, P., A. Fer and S. Renaudin. **1993**. Identification of the main osmotically active solutes in the unstressed and water-stressed root-hemiparasitic angiosperm *Thesium humile* and its host *Triticum vulgare*. Australian Journal of Plant Physiology **20: 223-230**.
- Simier, P., S. Renaudin and A. Fer. **1994**. Characteristics of the mannitol pathway in a root hemiparasitic species, *Thesium humile* Vahl. (Santalaceae). Journal of Plant Physiology **143: 33-38**.
- Yan, Z. **1993**. Resistance to haustorial development of two mistletoes *Amyema preissii* (Miq.) Tieghem and *Lysiana exocarpa* (Behr.) Tieghem ssp. *exocarpi* (Loranthaceae) on host and non-host species. International Journal of Plant Sciences (previously Botanical Gazette) **154: 386-394**. (Describes various types of resistance, in phloem and in xylem).
- Webb, M., A. Togola and D. Traore. **1993**. Effect of intercropping pearl millet with cowpea on the density of emerged *Striga hermonthica* in Mali. In: Proceedings Brighton Crop Protection Conference Weeds - **1993**, pp. **925-930**. (Cowpea intercrop reduced soil temperature and *Striga* density when planted in-row at time of cereal planting, but not when delayed).
- Wiesner, E. **1993**. Baeume und blumen in zentralen Alborz gebirge (Nord-Iran). Der Palmengarten (Munich) **2/93: 82-88**. [Figure 11 is a full color picture of *Phelypaea coccinea* (Orobanchaceae)].

HAUSTORIUM is edited by L. J. Musselman, Parasitic Plant Laboratory, Department of Biological Sciences, Old Dominion University, Norfolk, Virginia **23529-0266** USA, electronic mail LJM100F at ODUVM.CC.ODU.EDU, fax **804-683-5283** and C. Parker, C/O Long Ashton Research Station, University of Bristol, Bristol, **BS18 9AF**, ENGLAND, fax **(0275)394007**. It is prepared by John Musselman and is usually published twice yearly by Old Dominion University and funded by grant **DHR-5600-G-00-1021-00** from the Agency for International Development. Unsigned articles and literature reviews are by the editors. Send material for publication to either editor. Complete sets of HAUSTORIUM are available for **US\$30** postpaid. Make checks payable to Lytton J. Musselman and drawn on an American bank. Send requests for copies and sets to L. Musselman. If you have electronic mail, you may wish to receive your copies of HAUSTORIUM or the bibliographies in this manner. To do so, send a request via electronic mail to Lytton Musselman at the email address above. Issues **21-29** are available free via electronic mail.

Haustorium **28** was mailed **13 December 1993**
