

# HAUSTORIUM

## *Parasitic Plants Newsletter*

### Official Organ of the International Parasitic Seed Plant Research Group



February 1995 ..... Number 30

#### ● ● SIXTH INTERNATIONAL PARASITIC WEED SYMPOSIUM CORDOBA, SPAIN 16-18 APRIL 1996

Plan to attend the symposium in beautiful Cordoba, Spain where in addition to plenary sessions, papers, and posters there will be opportunity to see *Orobanche*, *Cuscuta*, and *Viscum* in the field. Like previous symposia, all groups of parasitic plants are included. For information contact:

Secretaria de "6th Parasitic Weed Symposium"  
Centro de Investigacion y Desarrollo Agrario  
hpartado 4240  
14080 Cordoba  
SPAIN  
Fax: (country code 34) 57 202721

#### ● BACTERIA THAT SCAVENGE GERMINATION STIMULANTS?

There is a microbiological principle stating no naturally occurring substance exists that cannot be broken down by a microorganism. *Striga* depends on the production of tiny seeds that are stimulated to germinate by chemical stimulants of which strigol is the best known. Strategy in the present research is to find a way to intervene into this process by interfering between the stimulant, whether strigol or one of its closely related analogs, and *Striga* seed. This project aims at finding a soil microorganism that could scavenge strigol as it exudes from the host root before it reaches the striga seed. Probability that such an organism can be found should be high because the microorganisms could have a number of other functions not confined to breaking down the probably recalcitrant heterocyclic strigol molecule. Simple techniques have been followed to test the hypothesis. Most of the work was carried out in sterile petri dishes with moistened filter papers on which sorghum

grains were grown (3-6 days) into seedlings which presumably release the strigol related substance sorgolactone on the filter paper. In the control experiments, preconditioned (10-12 days at 25 ° C) *Striga herrnonthica* seeds on glass fiber filter paper were placed in the petri dish with the sorghum seedlings, incubated at room temperature and the germinated seeds counted. The soil microorganisms to be tested for their ability to nullify the effect of the stimulant are grown on nutrient agar plates, the cell population suspended in distilled water and the suspension used in the test experiment. Here the bacterial suspension (inhibitor carrier) was added together with the *Striga* seeds to the petri dish carrying the sorghum seedlings. The isolate most commonly used was a mucoid, yellowish bacterium (probably a *Xanthomonas* sp.), originally isolated as a sorghum seed pathogen and was routinely propagated on sorghum seedlings in petri dishes. The addition of the bacterial suspension to the stimulant source reduced the ability to germinate the *Striga* seeds. About 65% of the *Striga* seeds germinated in the absence of the bacterial suspension (control) while only 8.4% germinated in the presence of the microorganisms (Treated.) In a supportive experiment, sorghum grains were grown in sterile sand in plastic cups for one week and watered with test bacterial suspension or with distilled water. Then the water was removed by a vacuum pump and used to stimulate *Striga* seeds. Controls gave 55.5% germination while test experiments gave 27.4% germination. A few pot experiments were also carried out in which sorghum seedlings were transferred from the petri dishes to large clay pots containing garden soil infested with *Striga* seeds at the rate of 0.08 grams per pot. The sorghum was grown for 44 days and plants watered with bacterial suspensions or with distilled water. Then, the plants were uprooted and the attachments of the *Striga* to the sorghum roots were counted. The results obtained showed that when bacterial suspensions were added to the soil the attach-

ments of striga to host roots averaged 36; in the controls, they averaged 114 attachments. We have also tried to gather some preliminary information on the nature of the inhibitor. In one experiment, we separated the liquid from the cell mass in the suspension, using a 0.45 micron bacterial filter, and tested the ability of each to inhibit the germination stimulant. The effective ingredient resided in the filtrate not in the cell mass. Heat treatment destroyed the ability of the inhibitor to have any effect on the stimulant suggesting that the inhibitor is a volatile substance or a protein that is denatured by heat. From a biotechnological view point, once a suitable organism has been obtained and the mechanism of its action on the stimulant has been elucidated, the useful ingredient can be obtained by growing the organism in a suitable growth medium in bioreactors. If the cells themselves are to be mixed with the seeds of sorghum before sowing as a dressing, then they can be provided in dry powder form like any other commercial microbial preparation. Obviously microbes that are pathogenic (e.g., the strain used here) cannot be used. If a chemical produced by the organism is the effective ingredient, then it could be produced in pure form. Both preparations should pose no environmental problems. The technology needed for the production of large scale biomass or chemical is available today. The yields can be improved through modern techniques of process optimization, and the microbial strains could be improved using recombinant DNA technology. It may even be possible to transfer the genes concerned from the microorganism to the host plant itself to make it intrinsically resistant to *Striga*. We believe that this research could open an area of useful research in weed control as the technique could equally well be applied to the control of similar parasitic weeds.

Hamid Ahmed Dirar, Faculty of Agriculture, Omar Mukhtar University, Al Beida, Libya

---

### ● SEX RATIO IN MISTLETOES

There are many dioecious mistletoes from the new and the old world. The sex ratio of many dioecious plants often deviates significantly from 1:1 with tendency towards a male bias. Earlier sex ratio surveys on mistletoes have reported a female-bias. *Viscum album* populations have a strongly female biased sex ratio in natural populations, but in other European and Asian species of *Viscum* sex ratios not differing from 1:1 have been found. Sex ratios were at unity in

most dioecious African species of *Viscum*, but female biased ratios as low as 0.52 and male biased ratios as high as high 1.40 may occur in some species. Female biased sex ratios were found over all populations of *Phoradendron tomentosum* in Central Texas by Nixon and Todzia. Recently Marshall et al. stated that, unique among the mistletoes studied to date, the sex ratio of *Phoradendron juniperinum* populations is significantly male biased. In populations of *Loranthus europaeus*, parasitizing oaks (*Quercus cerris* and *Q. petraea*), I found male bias. Male plants prevailed: they formed 69.2% of all living plants. This male biased sex ratio was in relation to the woodland coenopopulation of *Loranthus*. Biological and ecological causes of the variations in sex ratios in mistletoes are not known. Barlow found that in dioecious species of *Viscum*, males are normally heterozygous for sex associated chromosomal translocations, and in *V. album* male plants usually form a ring of eight to ten chromosomes at meiosis. Nixon and Todzia found that only the trees with one mistletoe exhibited a sex ratio near 1:1. Trees with two or more mistletoes generally showed female biased ratios. The authors suggested that a general trend of increased within tree female bias is associated with higher number of mistletoe per tree. But *Loranthus europaeus* showed similar sex ratios in both the woodland and within host tree populations. We need more information about sex ratios in several mistletoes species and in several environmental conditions. In mistletoe population dynamic studies it is important to accept the dioecy and sex ratio bias.

Pavol Elias, Institute of Botany, Slovak Academy of Sciences, Dubravská 14, SK-842 23 Bratislava, Slovakia

---

### ● PROCEEDINGS OF THIRD OROBANCHE SYMPOSIUM

The proceedings can be obtained from: KIT Press, Royal Tropical Institute, Mauritskade 63, 1092 AD Amsterdam, NETHERLANDS. The cost is 55 US\$ + \$12 for mailing surface mail and \$24 for air. Payment is to be made upon receipt of the volume. When ordering, indicate mode of shipment. (See literature section below under Pieterse etc for complete citation. See HAUSTORIUM 29 for a report on the symposium.)

---

### ● LITERATURE

- Abbasher Awad Abbasher. 1994. Microorganisms associated with *Striga hermonthica* and possibilities of their utilization as biological control agents. *PLITS* 12 (1): 144 p. (28 fungal species recorded in Sudan of which *Fusarium nygamai* and *F. semitectum* var. *majus* show potential for development as mycoherbicides.)
- Abu-Irmaileh, B. E. 1994. Nitrogen reduces branched broomrape (*Orobancha ramosa*) seed germination. *Weed Science* 42: 57-60. (Laboratory study showing that N 20-100 mg/l, as ammonium nitrate, reduced germination of *O. ramosa* when host seedlings and parasite seed were continuously exposed for 30-35 days.)
- Alabi, M. O., D. K. Berner and G. O. Olanyan. 1994. Characterization of soybean cultivars for *Striga hermonthica* control. Abstract 671. *Phytopathology* 84: 1151.
- Andrews, D. J. and P. J. Bramel-Cox. 1993. Breeding cultivars for sustainable crop production in low-input dryland agriculture in the tropics. pp 211-223 in: Buxton, D. R. et al. (Eds) *International Crop Science 1. International Crop Science Congress, Ames, 1992. Crop Science Society of America.* (Including consideration of sorghum resistance and yield loss due to *Striga* spp.)
- Andary, C., R. Wylde, L. Maury, A. Heitz, A. Dubourg, and S. Nishibe. 1994. X-ray analysis and extended NMR study of orapoxide. *Phytochemistry* 37(3): 855-857.
- Ariga, E. S., D. K. Berner and J. Chegwa. 1994. Effects of previous season cotton and cowpea on *Striga hermonthica* parasitism on maize. Abstract 670. *Phytopathology* 84: 1151.
- Aronne, G., C. C. Wilcox and P. Pizzolongo. 1993. Pollination biology and sexual differentiation of *Osyris alba* (Santalaceae) in the Mediterranean region. *Plant Systematics and Evolution* 188 (2): 1-16.
- Assefa Admasu and Endale Berhe. 1994. *Orobancha crenata* a potential threat to food legumes in Ethiopia. *EWSS Newsletter (Ethiopian Weeds Science Society)* 2(1): 1-2. (The first record of this species in Ethiopia, at two sites near Dessie - a threat to the important faba bean crop.)
- Atkinson, K., editor. 1994. IITA Annual Report 1993. International Institute of Tropical Agriculture. (Includes section on *Striga* research accomplishments.)
- Babiker, A. G. T., T. Cai, G. Ejeta, L. G. Butler and W. R. Woodson. 1994. Enhancement of ethylene biosynthesis and germination with thidiazuron and some selected auxins in *Striga asiatica* seeds. *Physiologia Plantarum* 91: 529-536. (confirming a synergistic interaction between thidiazuron and auxins.)
- Banerjee, K., D. C. Khatua and N. Mukherjee. 1993. Some new hosts of *Cuscuta* sp. *Indian Forester* 119: 760-761. (Recording occurrence on a range of forest tree species, leading to death of *Millingtonia hortensis*.)
- Berner, D. K., A. E. Awad and E. I. Aigbolchan. 1994. Potential of imazayuin seed treatment for control of *Striga gesnerioides* and *Alectra vogelii* in cowpea (*Vigna unguiculata*.) *Plant Disease* 78: 18-23. (Soaking cowpea seeds in imazayuin, e.g. 3.6 mg/l for 5 minutes before sowing, almost prevents parasite development.)
- Butler, L. G. 1995. Chemical communication between the parasitic weed *Striga* and its crop host. A new dimension in allelochemistry. pp. 158-168 in Dakshini, K. M. M. and F. A. Einhellig, editors. *Allelopathy. Organisms, Processes, and Applications.* Washington: American Chemical Society. ACS Symposium Series 582. (A helpful summary of chemical signals in the *Striga* life cycle.)
- Canevari, M. and E. Sieckert. 1993. Timing and rate of MON 13200 for dodder control in alfalfa. In: Lym, R. G. (Ed.) *Proceedings, Western Society of Weed Science* 46: 20-21. (Thiazopyr at 1 lb/ac applied to established alfalfa in January provided good control of *Cuscuta* through to August.)
- Carsky, R. J., L. Singh and R. Ndikawa. 1994. Suppression of *Striga hermonthica* on sorghum using a cowpea intercrop. *Experimental Agriculture* 30: 349-358. (More evidence for the possible value of cowpea interplanted within the row in reducing *Striga* density and seed production.)
- Carsky, R. J., L. Singh and R. Ndikawa. 1994. Effect of herbicide and handweeding on current and subsequent season *Striga hermonthica* density on sorghum. *International Journal of Pest Management* 40: 111-116. (No significant effect from 2 seasons of suppression, but initial infestation very high - >100/m<sup>2</sup>.)
- Carter, R. J. and D. A. Cooke. 1994. Broomrape (*Orobancha* spp.) is a threat to broadleaved crops in Australia. *Plant Protection Quarterly* 9: 61-63. (Including a useful key for separation of the main weedy species, not yet established in Australia, from the native *Orobancha cernua* var. *australiana*.)
- Cechin, I. and M. C. Press. 1994. Influence of nitrogen on growth and photosynthesis of a C<sub>3</sub> plant, *Oryza sativa* infected with the root hemiparasite,

- Striga hermonthica*. Journal of Experimental Botany 45: 925-930. (Photosynthesis reduced by 30 and 44% at high and low N respectively.)
- Cechin, I. and M. C. Press. 1994. The influence of nitrogen on growth and photosynthesis of sorghum infected with *Striga hermonthica* from different provenances. Weed Research 34: 289-298.
- Chowdhury, Al-Mamum and A. Ahmed. 1993. Histopathological studies of *Striga* spp. with sugarcane roots. Journal of Mycopathological Research 31(1): 1-4. (Observations on *S. asiatica* and *S. densiflora*.)
- Condon, J. and J. Kuijt. 1994. Anatomy and ultrastructure of the primary endophyte of *Ileostylus micranthus* (Loranthaceae.) International Journal of Plant Science 155(3): 350-364.
- Cubero, J. I., S. Khalil, and M. T. Moreno. 1993. Problems and prospects of selecting for *Orobanchae* resistance in cool-season legumes. pp. 167-177 in Singh, K. B. and Saxena, M. C. (Eds) Breeding for Stress Tolerance in Cool-season Food Legumes. John Wiley.
- Dawoud, D. A. and J. Sauerborn. 1994. Impact of drought stress and temperature on the parasitic weeds *Striga hermonthica* and *Alectra vogelii* in their early growth stages. Experimental Agriculture 30: 249-257.
- Dawson, J., Musselman, L. J., Doerr, I. and P. Wolswinkel. 1994. Biology and Control of *Cuscuta*. Reviews of Weed Science 6: 265-317. (A comprehensive survey of virtually every aspect of dodders: ecology, floral biology, taxonomy, distribution, ethnobotany, physiology, haustorial structure, control and more with extensive bibliography and numerous illustrations.)
- Dembele, B., A. Raynal-Roques, G. Salle, and C. Tuquet. 1994. Plantes Parasites des Cultures et des Essences Forestieres au Sahel. CTA/John Libbey Eurotext. 43 pp. (A beautifully illustrated account of a wide range of parasitic species in West Africa.)
- Dieringer, G. and C. R. Werth. 1994. Population genetic structure of *Agalinis strictifolia* (Scrophulariaceae.) Bulletin of the Torrey Botanical Club 121(2): 148-153.
- Erskine, W. and M. C. Saxena. 1993. Problems and prospects of stress resistance breeding in lentil. In: Singh, K. B. and Saxena, M. C. (Eds) Breeding for Stress Tolerance in Cool-season Food Legumes. John Wiley. pp. 51-62.
- Ersline, W., M. Tufail, A. Russel, M. C. Tyagi, M. M. Rahman and M. C. Saxena. 1994. Current and future strategies in breeding lentil for resistance to biotic and abiotic stresses. Euphratica 73: 127-135. (Noting the lack of sources of resistance to *Orobanchae* spp. in lentil.)
- Fer, A., J. C. Painset and L. Rey. 1992. [Characteristics of mineral nutrition in some types of parasitic plants (*Cuscuta*, *Orobanchae*, *Thesium*) and their development on crops.] Bulletin de la Societe Botanique de France, Actualites Botaniques 139: 111-121. (Observations on K levels in host and parasite suggest that *Cuscuta* and *Orobanchae* obtain nutrients mainly from the host phloem, while *Thesium* obtains them from host xylem.)
- Fer, A., N. Russo, P. Simier, M-C. Arnaud and P. Thalouarn. 1994. Physiological changes in a root hemiparasitic angiosperm, *Thesium humile* (Santalaceae) before and after attack. Journal of Plant Physiology 143: 704-710. (Growth and photosynthetic rate in unattached parasite declines after 2 weeks, perhaps associated with high N/low P uptake by parasite roots. Initial photosynthesis comparable with that of wheat.)
- Fernandez-Rivera. S. and D. E. Hess. 1994. Chemical composition of *Striga* herbage and its intake and digestion by sheep. Abstract 675. Phytopathology 84: 1151.
- Filip, G. M., J. J. Colbert, C. G. Shaw, P. Hessburg and K. P. Hosman. 1993. Influence of dwarf mistletoe and western spruce budworm on growth and mortality of Douglas-fir in unmanaged stands. Forest Science 39: 465-477. (Results showed serious effects from both pests but no synergistic interaction between them.)
- Frischmuth, K., A. Marx, T. Petrowitsch, U. Wagner, K. Koerner, S. Zimmerman, H. Meuer, W. S. Sheldrick, and P. Welzel. 1994. Strigol synthetic studies: The first synthetic approach that allows control of C-2' configuration. Tetrahedron Letters 35(28): 4973-4976.
- Garcia-Torres, L., F. Lopez-Granados and M. Castejon-Munoz. 1994. Pre-emergence herbicides for the control of broomrape (*Orobanchae cernua* Loefl.) in sunflower (*Helianthus annuus* L.) Weed Research 34: 387-402. (Reporting highly selective control from imazethapyr, imazapyr and chlorsulfuron.)
- Gomes, A. L. and G. W. Fernandes. 1994. Influence of parasitism by *Pilostyles ingae* (Rafflesiaceae) on its host plant *Mimosa nagiurei* (Leguminosae.) Annals of Botany 74: 205-208.
- Grimble, D. G. and R. C. Beckwith 1993. Temporal changes in presence of late instar *Mitoura spinetorum* (Lyceaniidae) in eastern Oregon. Journal of the Lepidopterists' Society 47: 329-330. (On

- Arceuthobium campylopodum* on *Pinus ponderosa*.)
- Haberhausen, G. and K. Zetsche. 1994. Functional loss of all ndh genes in an otherwise relatively unaltered plastid genome of the holoparasitic flowering plant *Cuscuta reflexa*. *Plant Molecular Biology* 24: 217-222.
- Hanounik, S. B., G. J. Jellis and M. M. Hussein. 1993. Screening for disease resistance in faba bean. pp 97-106 in Singh, K. B. and Saxena, M. C. (Eds) *Breeding for Stress Tolerance in Cool-season Food Legumes*. John Wiley. (Including reference to *Orobanche crenata*.)
- Hawksworth, F. G. and D. W. Johnson. 1993. You can save your trees from dwarf mistletoe. USDA Forest Service Rocky Mountain Forest and Range Experiment Station. General Technical Report RM-225. 10 pp. (Control methods include felling, pruning, spraying with ethephon every 3-5 years, improving tree vigor, planting resistant tree species.)
- Hess, D. E. 1994. Crop-specific strains of *Striga hermonthica* in Niger. Abstract 674. *Phytopathology* 84: 1151.
- Hsiao, S., J. D. Mauseth, and C. Peng. 1995. Composite bundles, the host/parasite interface in the holoparasitic angiosperms *Langsdorffia* and *Balanophora* (Balanophoraceae.) *American Journal of Botany* 82(1): 81-91. (Composite bundles are the host parasite interface in these plants. Transfer cells are present and may represent the main pathway of interchange between host and parasite.)
- INTSORMIL (Sorghum and Millet Collaborative Research Support Program). 1994. Bibliography. INTSORMIL publication 94-5. (Includes a separate section with 51 references to work on *Striga*.)
- Jeschke, W. D., P. Baumel, N. Raete, F-C. Czygan and P. Protsch. 1994. Modelling the flows and partitioning of carbon and nitrogen in the holoparasitic *Cuscuta reflexa* Roxb. and its host *Lupinus albus* L. II Flows between host and parasite and within the host. *Journal of Experimental Botany* 45: 801-812. (Showing severe drain on host N as well as carbohydrate.)
- Joel, D. M. and D. Losner-Goshen. 1994. The attachment organ of the parasitic angiosperms *Orobanche cumana* and *O. aegyptiaca* and its development. *Canadian Journal of Botany* 72: 564-574.
- Khallide, R., S. P. S. Beniwal, Z. Fatemi and M. C. Saxena. 1993. Imazethapyr can control dodder (*Cuscuta* spp.) infestation in faba bean. FABIS Newsletter 33: 30-32. (Doses of 75 g ai/ha pre- or 20 g ai/ha post-emergence intended for *Orobanche* control found to be highly selective v. *Cuscuta*.)
- Rim, S. K. 1994. Genetics of maize tolerance of *Striga hermonthica*. *Crop Science* 34: 900-907. (Concluding that the genetic control of tolerance is polygenic and the inheritance quantitative.)
- Kim, S. K., A. Y. Akintunde and P. Walker. 1994. Response of maize, sorghum and millet host plants to infestation by *Striga hermonthica*. *Crop Protection* 13: 582-590.
- Kirk, A. A. 1994. A fungal pathogen with potential for control of *Striga hermonthica* (Scrophulariaceae.) *Entomophaga* 38: 459-460. (*Fusarium equiseti* affected 50% of *Striga* in Sunjikai district, Kordofan. Sudan and prevented seeding in most plants.)
- Kondap, S. M. and R. M. Kumar. 1993. Management of *Cuscuta* in croplands and fallows. pp 407-411 in *Integrated Weed Management for Sustainable Agriculture*. Proceedings of an Indian Society of Weed Science International Symposium, Hisar, 1993. Volume I.
- Krishnamurty, G. V. G., M. S. Chari and K. Nagarajan. 1994. Broomrape (*Orobanche cernua* Loefl.) on tobacco. Central Tobacco Research Institute, Rajahmundry - 533105, India. 35 pp. (A useful, well-produced summary of biological and control studies in India.)
- Kuijt, J. 1994. Typification of the names of New World mistletoe taxa (Loranthaceae and Viscaceae) described by Martius and Eichler. *Taxon* 43: 187-199.
- Kuijt, J. 1994. A hyperparasite mistaken for an inflorescence: The identify and typification of *Phoradendron rusbyi* Britton (Viscaceae.) *Novon* 4: 116-118.
- Kumar, R. M., S. M. Kondap, N. Madhav Rao and A. Ramakrishna. 1993. An anatomical study on host specificity of a parasitic weed - *Cuscuta*. pp 61-63 in: *Integrated Weed Management for Sustainable Agriculture*. Proceedings of the Indian Society of Weed Science International Symposium, Hisar, 1993. Vol. 2.
- Kuppers, M., B. I. L. Kuppers and A. G. Swan. 1993. Leaf conductances and xylem pressures of the host/mistletoe pair *Eucalyptus behriana* F. Muell. and *Amyema miquelii* (Lehm. ex Miq.) Tiegh. at permanently low plant water status in the field. *Trees: Structure and Function* 8: 110-114. (Transpiration was always greater in the host than in the parasite.)

- Lahrada, R., J. C. Caselev and C. Parker. 1994. Weed Management for Developing Countries. FAO Plant Production and Protection Paper 120. FAO, Rome. 384 pp. (Includes sections on *Cuscuta* by Li, *Striga* by Musselman and *Orobancha* by Sauerborn. Available on request from the senior author, Plant Protection Service, FAO, Rome 00100, Italy.)
- Leake, J. R. 1994. Tansley Review No 69. The biology of myco-heterotrophic ('saprophytic') plants. *New Phytologist* 127: 171-216. (A comprehensive review relating to 400 species in 87 genera, discussing among much else whether 'saprophytic' plants are symbiotic with, or parasitic upon, their associated fungi.)
- McKinnel, F. H. (ed.) 1993. Sandalwood in the Pacific Region. Australian Council for International Agricultural Research. 43 pp. (Surveying biodiversity, cultural practices and uses for timber, aromatic oil and fruit.)
- Marshall, J. D., J. R. Ehleringer, E. D. Schulze and G. Farquhar. 1994. Carbon isotope composition, gas exchange and hypertrophy in Australian mistletoes. *Functional Ecology* 8: 237-241.
- Mauseth, J. D. 1993. Cacti, mistletoes and desert survival. *Cactus and Succulent Journal* 65: 172-175. (Concerning *Tristerix aphyllus* on cacti in Chile.)
- Mbuga, A. M. and A. T. Obilana. 1993. Distribution and host specificity of *Striga asiatica* and *S. hermonthica* on cereals in Tanzania - a preliminary study. *International Journal of Pest Management* 39: 449-451.
- Moors, A., D. K. Berner and P. Van Damme. 1994. Identification of nonhosts for control of *Striga gesnerioides*. Abstract 672. *Phytopathology* 84: 1151.
- Nair, A. G. R. and G. Thirupurasundari. 1992. Coumarins and flavonoids from *Cuscuta reflexa* parasitic on *Bougainvillea spectabilis*. *Fitoterapia* 63: 381-382.
- Ndirpaya, Y. D., D. K. Berner and L. J. Musselman. 1994. Germination requirements of *Striga aspera*. Abstract 673. *Phytopathology* 84: 1151.
- Nickrent, D. L., K. P. Schuette and E. M. Starr. 1994. A molecular phylogeny of *Arceuthobium* (Viscaceae) based on nuclear ribosomal DNA internal transcribed spacer sequences. *American Journal of Botany* 81: 1149-1160.
- Overton, J. M. 1994. Dispersal and infection in mistletoe in populations. *Journal of Ecology* 82: 711-723. (*Phryglanthus sonorae* (Loranthaceae) in Baja California shows little or no host preference.)
- Panda, M. M. and N. K. Choudhury. 1993. Effects of nutrients on chlorophyll content and photochemical activities by isolated chloroplasts of *Cuscuta reflexa*. *Plant Physiology and Biochemistry* (New Delhi) 19: 6-10.
- Pate, J. S., G. Woodall, W. D. Jeschke and G. R. Stewart. 1994. Root xylem transport of amino acids in the root hemiparasitic shrub *Oxalophyllanthi* (Labill.) R. Br. (Olacaceae) and its multiple hosts. *Plant, Cell and Environment* 17: 1263-1273. [Amino acids in the parasite occasionally reflect those of the host but 'pronounced metabolic transformations' are demonstrated. Some observations also on *Exocarpos spartens* (Santalaceae) which was parasitized by, but did not parasitize, *Oxal.*]
- Pazy, B. and U. Plitmann. 1994. Holocentric chromosome behaviour in *Cuscuta* (Cuscutaceae.) *Plant Systematics and Evolution* 191: 105-109.
- Pieterse, A. H., J. A. C. Verkleij and S. J. ter Borg (Eds.) 1994. Biology and Management of *Orobancha*. Proceedings of the Third International Workshop on *Orobancha* and related *Striga* Research. Royal Tropical Institute, The Netherlands. 736 pp. (Containing 96 papers, which will not be separately listed in *Haustorium* - see separate announcement for further information.)
- Porta-Puglia, A., K. B. Singh and A. Infantino. 1993. Strategies for multiple-stress resistance breeding in cool-season food legumes. pp. 411-427 in K. B. and Saxena, M. C. (Eds) *Breeding for Stress Tolerance in Cool-season Food Legumes*. John Wiley.
- Prance, G. T. and P.H. Raven, editors. 1995. *The World of Plants*. Weekly Encyclopedia 40(1/22): 97-128. Tokyo: Asahi Shimbun Publishers. (Popular treatment with spectacular photographs of the most bizarre of all angiosperm families: the Rafflesiaceae, Mistrastemonaceae, Hydnoraceae, Balanophoraceae, Eremolepidaceae, Viscaceae, Loranthaceae, Misodendraceae, Santalaceae, Opiliaceae, Olacaceae, Diptentodontaceae, and Medusandraceae by leading students of each family. In Japanese but with Latin names on photographs.)
- Prather, L. A. and R. J. Tyrl. 1993. The biology of *Cuscuta attenuata* Waterfall. *Proceedings of the Oklahoma Academy of Science* 73: 7-13. (Unlike most species of dodder, *C. attenuata* has a restricted host range, preferring *Iva annua*; Asteraceae.)
- Press, M. C., A. N. Parsons, A. W. Maclay, C. A. Vincent, V. Cochrane and W. E. Seel. 1993. Gas

- exchange characteristics and nitrogen relations of two Mediterranean root hemiparasites: *Bartsia trixago* and *Parentucellia tiscosa*. *Oecologia* 95: 145-151.
- Press, M. C. and J. B. Whittaker. 1993. Exploitation of the xylem stream by parasitic organisms. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences* 341: 101-111. (Relating especially to *Striga asiatica* and *Viscum album*.)
- Ramlan, M. F. 1994. Environment and the host-parasite interactions between *Striga hermonthica* and sorghum. PhD Thesis. York University, UK. 161 pp.
- Rao, P. N. and K. K. Neelai. 1993. Diversity in the host range of China dodder. pp. 1-6 in K. Udaiyan. K. Janardhanan, S. Manian and V. R. K. Reddy (Eds) Proceedings of the National Seminar on biodiversity : strategies for conservation and future challenges. Coimbatore, 1993. (Listing a wide range of hosts of *Cuscuta chinensis* in India, including the crops pigeon pea, sunn hemp. *Phaseolus trilobus* and *Vigna mungo*.)
- Sharma, S. B., D. H. Smith, P. Subrahmanyam, M. J. Freire, T. Faria, D. R. Butler and G. V. Ranga Rao. 1994. Parasitism of groundnut by *Striga* sp. in Mozambique. *International Arachis Newsletter* 14: 24-25. (Confirming the parasitism of groundnut by an identified *Striga* sp., likely *S. gesnerioides*.)
- Riches, C. R. 1994. Variability of parasitic weeds and their hosts: implications for witchweed management in Africa. In: *The Impact of Genetic Variation on Sustainable Agriculture. Aspects of Applied Biology* 39. Association of Applied Biology. pp. 145-154.
- Riches, C. R., L. J. Shaxson, J. W. M. Logan and D. C. Munthali. 1993. Insect and parasitic weed problems in southern Malawi and the use of farmer knowledge in the design of control measures. *Network Paper - Agricultural Administration (Research and Extension) Network* 42: 1-17. ODA, London.
- Runge-Metzger, A. 1993. The economics of *Striga* control in different farming systems in northern Ghana. Report PN 87.3540.9-01.200, Supra-regional Project, Ecology and Management of Parasitic Weeds. GTZ-UH. 51 pp. (A thoughtful analysis of various socio-economic factors, emphasizing the need for more long-term studies and data.)
- Saghir, A. R. 1994. An overview of a new technique in *Orobanche* control. *Plant Protection Quarterly* 9: 77-80. (Referring to the use of synthetic strigol analogs.)
- Shaw, C. G. and D. Phanichapol. 1993. First report of *Dendrophthoe falcata* on *Pinus kesiya*. *Plant Disease* 77: 847.
- Shaxon, L. J., C. R. Riches and J. H. Seyani. 1993. Incorporating farmer knowledge in the design of weed control strategies for smallholders. *Proceedings, Brighton Crop Protection Conference, Weeds*. 1993. Vol. 3. pp. 1149-1154. (With particular reference to *Striga* spp. in Malawi.)
- Singh, A. and Singh, M. 1993. Cell wall degrading enzymes in *Orobanche aegyptiaca* and its host *Brassica campestris*. *Physiologia Plantarum* 89: 177-181. (Suggesting cellulase and polygalacturonase to be the main enzymes involved in haustorial penetration.)
- Smith, L. C. 1993. Studies on *Zea* mays infected with *Striga hermonthica*, with particular reference to photosynthesis. PhD Thesis, University of London. 170 pp. (*Striga* infection resulted in reduction in photosynthesis, perhaps via changes in hormone levels and resultant changes in cell wall structure.)
- Smith, R. B., E. F. Wass and M. D. Meagher. 1993. Evidence of resistance to hemlock dwarf mistletoe (*Arceuthobium tsugense*) in western hemlock (*Tsuga heterophylla*) clones. *European Journal of Forest Pathology* 23: 163-170.
- Sontakey, P. Y., W. M. Dabre, R. D. Deotale, V. K. Mohod and D. R. Kene. 1992. Correlation among morpho-physiological and biochemical constituents in *Striga* resistant and susceptible cultivars of sorghum. *Journal of Soils and Crops* 2(1): 32-34. (Showing positive correlation between *Striga*-resistance and hydrocyanic acid levels in leaves at 30 days.)
- Soria, S., D. Cadahia and A. Munoz. 1993. (The genus *Carulaspis* Mac. Gillivray, 1921 (Homoptera, Diaspididae) in the gardens of Patrimonio Nacional.) *Boletín de Sanidad Vegetal, Plagas* 19: 273-284. (Occurs on *Viscum album* in Spain.)
- Srivastava, S., A. Nigohjkar and A. Kumar. 1994. Multiple forms of pectin methylesterase from *Cuscuta reflexa* filaments. *Phytochemistry* 37: 1233-1236.
- Suzuki, Y., Yun-Hui Zhang, N. Murofushi and Y. Takeuchi. 1994. Endogenous gibberellins in clover broomrape, a parasitic plant and its host, clover: dependency of the parasite on the host for gibberellin production. *Journal of Plant Growth Regulation* 13: 63-67. (The range of gibberellins in *O. minor* is similar to those in the host, sug-

- gesting the host as the main source; however, some are found only in the parasite suggesting some ability for gibberellin synthesis in the parasite.)
- Sweeney, E. C., R. A. Palmer and U. Pfüller. 1993. Crystallization of the ribosome inactivating protein ML1 from *Viscum album* (mistletoe) complexed with beta-D-galactose, *Journal of Molecular Biology* 234: 1279-1281.
- ter Borg, S. J., A. Willemsen, S. A. Khalil, H. A. Saber, J. A. C. Verkleij and A. H. Pieterse. 1994. Field study of the interaction between *Orobanche crenata* Forssk. and some new lines of *Vicia faba* L. in Egypt. *Crop Protection* 13: 611-616. (Confirming partial resistance in Giza 402 and some derivatives from it.)
- Tsuda, T., A. Sugaya, Y. Z. Liu, K. Kathoh, H. Tanaka, H. Kawazura, E. Sugaya, E. Kusai, and M. Kuhno. 1994. Radical scavenger effect of *Boschniakia rossica*. *Journal of Ethnopharmacology* 41(1/2): 83-90.
- Uhlich, H. 1994. [*Orobanche baumanniorum* var. *longebracteata* (Orobanchaceae) - a new variety from Greece.] *Willdenowia* 24: 49-51. (In German.)
- van Hezewijk, M. J., K-H. Linke, F. Lopez-Granados, O. A. Al-Menoufi, L. Garcia-Torres, M. C. Saxena, J. A. C. Verkleij and A. H. Pieterse. 1994. Seasonal changes in germination response of buried seeds of *Orobanche crenata* Forsk. *Weed Research* 34: 369-376. (Confirming a pronounced natural cycle of germinability, during winter months only.)
- Wolf, S. J. and M. P. Timko. 1994. Characterization of actin-gene family members and their expression during development in witchweed (*Striga asiatica* L.) *Planta* 192: 61-68.
- Yahara, S., H. Domoto, C. Sugimura, T. Nohara, Y. Niiho, Y. Nalcjima and H. Ito. 1994. An alltaloid and two lignans from *Cuscuta chinensis*. *Phytochemistry* 37: 1755-1757. (Isolated from a preparation of dried fruits used in China as part of a drug complex to cure impotence and as a tonic.)

---

HAUSTORIUM is edited by L. J. Musselman, Parasitic Plant Laboratory, Department of Biological Sciences, Old Dominion University, Norfolk, Virginia 23529-0266 USA, electronic mail LJM100F at ODUVM.CC.ODU.EDU, fax 804-683-5283 and C. Parker, c/o Long Ashton Research Station, University of Bristol, Bristol, BS18 9AF, ENGLAND, fax (1275) 394007. It is prepared by John Musselman and is usually published twice yearly by Old Dominion University and funded by grant DHR-5600-G-00-1021-00 from the Agency for International Development. Unsigned articles and literature reviews are by the editors. Send material for publication to either editor. Complete sets of HAUSTORIUM are available for US\$30 postpaid. Make checks payable to Lytton J. Musselman and drawn on an American bank. To receive your copies of HAUSTORIUM or the bibliographies via electronic mail, send a message to the email address above. Issues 21-30 are available electronically.

---

Haustorium 29 was mailed 2 June 1994

---